



Aflatoxin B1 Affects Kenyan Markets: How It Can be Managed

Solomon Omwoma Lugasi^{1*}

¹Department of Physical Sciences, Jaramogi Oginga Odinga University of Science and Technology, Kenya.

Author's contribution

The sole author designed, analysed, interpreted and prepared the manuscript.

Article Information

DOI: 10.9734/AIR/2019/v20i630175

Editor(s):

(1) Dr. Paola Deligios, Department of Agriculture, University of Sassari, Italy.

Reviewers:

(1) Pucci Graciela, National University of Patagonia San Juan Bosco (Universidad Nacional de la Patagonia San Juan Bosco), Argentina.

(2) Silvia Denise Peña Betancourt, Universidad Autónoma Metropolitana, Mexico.
Complete Peer review History: <http://www.sdiarticle4.com/review-history/52942>

Mini-review Article

Received 28 September 2019

Accepted 02 December 2019

Published 14 February 2020

ABSTRACT

Aflatoxin B1 is a toxin produced as secondary fungal metabolites by the fungus *Aspergillus*, particularly *A. flavus*. The toxin has significantly contaminated the food supply chain especially cereals in Kenya. Kenya Bureau of Standards (KEBS) has recently banned five maize flour brands, citing high aflatoxin levels. They also suspended seven peanut butter products and the permits of their parent companies over aflatoxin contamination. The huge losses encountered by these companies calls for concerted efforts to manage aflatoxin in cereals. Aflasafe, a natural product for controlling poisonous *A. flavus* in food crops, including maize is made from roasted sterile sorghum (usually colored blue using food color) coated with non-poison producing types of *A. flavus* native to Kenya. The product is broadcasted in the maize fields during flowering and after exposure to sufficient moisture, the friendly Aflasafe fungi grow out as green spores containing millions of spores that are eventually spread to the crop, carried by wind and insects in the manner that aflatoxin-producing fungi are spread.

Keywords: Aflatoxin; aflatoxin B1; *Aspergillus flavus*; aflasafe.

*Corresponding author: E-mail: solomwoma@yahoo.com;

1. INTRODUCTION

Aflatoxins are toxins produced as secondary fungal metabolites by the fungus *Aspergillus*, particularly *A. flavus*. The fungus produces aflatoxin B1 (AFB1), aflatoxin B2 (AFB2), aflatoxin G1 (AFG1), aflatoxin G2 (AFG2), aflatoxin M1 (AFM1) and aflatoxin M2 (AFM2) (Blake and Mustafa, 2019) (see Fig. 1). It has been proven epidemiologically and documented (IARC, 2012) that aflatoxins, especially AFB1 is a carcinogen. It causes hepatocellular carcinoma, suppresses growth, modulates the immune system and leads to malnutrition [1,2]. *A. flavus* is found in almost all soil types and it can contaminate the food chain either at flowering or post-harvest stages producing AFB1. High temperatures and humidity promote its growth [3].

AFB1 significantly contaminates the food supply chain especially cereals in Kenya. For instance, in 2013, farmers in Tana River County lost their maize harvest due to infections by aflatoxins [4]. The whole harvest at the Hola Agricultural Irrigation Scheme for maize of approximately 5,400 tones was affected by aflatoxins. The levels of aflatoxins were over the standard level 10 ppb. Maize reserves of about 207000 tons were also declared infected by aflatoxins in 2010 [4]. The trend continues in Kenya today making aflatoxin infection a threat to food supply chain. Most recently, Kenya Bureau of Standards (KEBS) banned five maize flour brands, citing high aflatoxin levels [5]. Their permits were suspended and the manufacturers instructed to discontinue manufacturing or offering for sale the

affected maize meal products. The five products were Dola by Kitui Four Mills, Kifaruru by Alpha Grain Limited, Starehe by Pan African Grain Millers, 210 Two Ten by Kenblest Limited and Jembe by Kensal Rise Limited [5]. Furthermore, KEBS also suspended seven peanut butter products and the permits of their parent companies over aflatoxin contamination. Among the products affected include Nuteez peanut butter, True nuts, Fressy, Supa meal, Sue's Naturals, Zesta and Nutty [6]. Definitely the aforementioned bans had serious financial implications.

2. HOW AFLATOXINS SPREADS

Although the fungus affects maize crop at the flowering stage, poor post-harvest practices including un-aerated storage and/or delays in reducing moisture content of the maize contribute immensely to aflatoxin infection/spread (see Fig. 2). Mechanized shredding of the maize that is not able to isolate affected maize also increases infections. It should be noted that aflatoxins infections are positively affected by dry climate. Therefore most of the irrigation schemes in dry areas for maize production in Kenya should consider aflatoxin infection as a threat. However, although sorting of cereals (maize) may help reduce fresh infections from *A. flavus* fungus whose infection is usually visible (see Fig. 3), AFB1 compound is colorless, odorless and tasteless. As such use of visual techniques to detect the compound can be misleading and should be avoided. Use of test kits that are readily available in the market should be encouraged.




Fig. 1. The structure of aflatoxins produced by the fungus *Aspergillus*, particularly *A. flavus*




Fig. 2. Health effects of aflatoxin B1. Adopted from [13]

3. PREVENTION OF AFLATOXIN B1 IN STORED CEREALS

Food security is part of the Kenyan Governments Agenda and as such appropriate post-harvest practices should be employed to reduce losses. The use of hermetic bags, cereal drying machines, grain cocoons or silos can reduce losses significantly. Furthermore, farmers can use the 'Alfasafe' technology to control *A. flavus* during the flowering stage [7].

4. HOW TO MANAGE AFLATOXINS B1 IN THE FIELD

Aflasafe is a natural product for controlling poisonous *A. flavus* in food crops, including maize. The product was developed by the Kenya Agricultural and Livestock Research Organisation (KALRO), jointly with the United States Department of Agriculture – Agriculture Research Service (USDA-ARS) and the International Institute of Tropical Agriculture (IITA) [8]. It is roasted sterile sorghum (usually colored blue using food color) coated with non-poison producing types of *A. flavus* native to Kenya [7]. The product is broadcasted in the maize fields during flowering and after exposure to sufficient moisture, the friendly Aflasafe fungi grow out as green spores containing millions of spores that are eventually spread to the crop, carried by wind and insects in the manner that aflatoxin-producing fungi are spread [9].




Fig. 3. Samples of maize affected with *A. flavus* fungus in Kenya

5. HOW TO MANAGE AFFECTED CEREALS

Other cases of aflatoxins infections in Kenyan cereals include peanut butter that was reported to contain aflatoxin, 24 ppb way above the recommended 10 ppb. As such, it is advisable for farmers to be sensitized so as to control aflatoxin during flowering and good post-harvest handling practices. Contaminated cereals beyond both human (<10 ppb) and animal (<300 ppb) [10] consumption maybe considered for generating ethanol for industrial use instead of total

destruction. Levels of AFB1 in such ethanol should be indicated as the toxin has been documented to show traces in the final finished fermented product [11]. However such ethanol can be used for general industrial use like in making printing inks, paints and coatings, screen wash and deicers for the automotive industry, etc [12].

6. HEALTH EFFECTS OF AFLATOXINS ON HUMAN AND ANIMALS (AFLATOXICOSIS)

Aflatoxicosis, a condition caused by aflatoxins, can be acute or chronic Fig. 2. Acute aflatoxicosis occurs upon consumption of moderate-high levels of aflatoxins. Symptoms include hemorrhage, acute liver damage, edema, alteration in digestion, absorption and/or metabolism of nutrients, and possibly death [13]. sorghum coated with non-poison producing types of *A. flavus*. After exposure to sufficient moisture, the friendly *Aflasafe* fungi grow out as green spores containing millions of spores that are eventually spread to the crop, carried by wind and insects in the manner that aflatoxin-producing fungi are spread.

The chronic aflatoxicosis results from ingestion of low to moderate levels of aflatoxins. Effects of chronic include impaired food conversions, retarded growth, congenital malformations, mutations in the genetic code, altering DNA, rearrangement of chromosome pieces, gain or loss of entire chromosomes, or changes within a gene and carcinogenic effect. The carcinogenic mechanisms have been identified such as the genotoxic effect where the electrophilic carcinogens alter genes through interaction with DNA and thus becoming a potential for DNA damage and the genotoxic carcinogens that are sometimes effective after a single exposure, can act in a cumulative manner, or act with other genotoxic carcinogens which affect the same organs [13,14]. More information on health effects of aflatoxins can be read from the following articles [13,15,16] etc.

7. CONCLUSION

It is important for companies to invest heavily in aflatoxin detection in cereals before buying and storing. Their storage should be robust including the use of hermetic bags, cereal drying machines, grain cocoons or silos. Farmers should also be advised to employ the *Aflasafe* technology in controlling field infections of aflatoxin B1. The technique involves the use of roasted sterile.

ACKNOWLEDGEMENT

Financial support from RSC Allan Ure Bursary fund of 2017 and Jaramogi Oginga Odinga University of Science and Technology are acknowledged.

COMPETING INTERESTS

Author has declared that no competing interests exist.

REFERENCES

1. IARC. Aflatoxins, IARC Monographs on the Evaluation of Carcinogenic Risks on Humans; 2012. Available:<https://monographs.iarc.fr/iarc-monographs-on-the-evaluation-of-carcinogenic-risks-to-humans-65/> 2012
2. Voth-Gaeddert LE, Stoker M, Torres O, Oerther DB. Association of aflatoxin exposure and height-for-age among young children in Guatemala. *Int J Environ Health Res.* 2018;28:280-292.
3. Iqbal SZ, Nisar S, Asi MR, Jinap S. Natural incidence of aflatoxins, ochratoxin A and zearalenone in chicken meat and eggs. *Food Control.* 2014;43:98-103.
4. Dominic O. Aflatoxin poses huge risk to food security. Standard Digital; 2019. Available:<https://www.standardmedia.co.ke/article/2001318991/aflatoxin-dents-kenya-s-food-security> (Accessed on 21/06/2019)
5. Nation. Kebs bans 5 maize flour brands over high aflatoxin levels; 2019. Available:<https://www.nation.co.ke/news/K-kebs-bans-5-maize-flour-brands/1056-5343060-eb9k78/index.html> (Accessed on 12th Nov. 2019)
6. Mireri J. Seven peanut butter brands suspended over aflatoxin contamination; 2019. Available:<https://www.standardmedia.co.ke/article/2001348078/seven-peanut-butter-brands-suspended-over-aflatoxin-contamination> (Accessed on 12th Nov. 2019)
7. Japhet R. *Aflasafe* protects your maize from deadly aflatoxin; 2017. Available:<http://farmbizafrica.com/pests/310-alfasafe-protects-your-maize-from-deadly-aflatoxin> (Accessed on 21/6/2019)
8. Gerald A. State approves new drug in aflatoxin war; 2015.

- Available:<https://www.businessdailyafrica.com/news/State-approves-new-drug-in-aflatoxin-war/539546-2787750-y0m0a6/index.html>
(Accessed on 11/25/2019)
9. Ranajit B, Matieyedou K. Taking Aflasafe from Science to Scale; 2018.
Available:<https://docs.lib.purdue.edu/cgi/viewcontent.cgi?article=1023&context=scal-eup>
(Accessed on 11/25/2019)
 10. Blake RR, Mustafa IS. Aflatoxin B1: A review on metabolism, toxicity, occurrence in food, occupational exposure and detoxification methods. *Food and Chemical Toxicology*. 2019;124:81-100.
 11. Tomonori I, Yasushi N, Atsuo U, Naoki M. Degradation of Aflatoxin B1 during the Fermentation of Alcoholic Beverages *Toxins*. 2013;5:1219-1229.
 12. IEA Industrial Ethanol; 2017.
Available:<http://www.industrial-ethanol.org/index.php?page=industrial-ethanol>
 13. Godfrey S, David K, Lubega A, Ogwal-Okeng J, William W, David B. Review of the Biological and Health Effects of Aflatoxins on Body Organs and Body Systems; 2013.
Available:<http://dx.doi.org/10.5772/51201>
 14. Wangikar PB, Dwivedi P, Sinha N, Sharma AK, Telang AG. Teratogenic effects in rabbits of simultaneous exposure to ochratoxin A and aflatoxin B1 with special reference to microscopic effects. *Toxicology*. 2005;215:37-47.
 15. Kensler TW, Roebuck BD, Wogan GN, Groopman JD. Aflatoxin: A 50 year odyssey of mechanistic and translational toxicology. *Toxicological Sciences*. 2010; 120:S28-S48.
 16. USAID. Aflatoxin: A Synthesis of the Research in Health, Agriculture and Trade; 2012.
Available:https://www.danya.com/portfolio/aflatoxin_report.pdf
(Accessed on 11/25/2019)

© 2019 Lugasi; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<http://www.sdiarticle4.com/review-history/52942>